

ISSN: 2321 9114
 AJEONP 2014; 1 (3): 16-18
 © 2014 AkiNik Publications
 Received 22-12-2013
 Accepted: 07-03-2014

William N. Setzer
 Department of Chemistry, University
 of Alabama in Huntsville, Huntsville,
 AL 35899, USA
 Email: wsetzer@chemistry.uah.edu

Chemical composition of the floral essential oil of *Tabernaemontana longipes* from Monteverde, Costa Rica

William N. Setzer

Abstract

The floral essential oil of *Tabernaemontana longipes* Donn. Sm. (Apocynaceae) from Monteverde, Costa Rica, was isolated by hydrodistillation and analyzed by gas chromatography – mass spectrometry. The floral oil was dominated by the monoterpene (*E*)- β -ocimene (30.0%), the sesquiterpene germacrene D (12.5%), and the normal alkane heneicosane (12.9%).

Keywords: *Tabernaemontana longipes*, essential oil composition, β -ocimene, germacrene D, heneicosane.

1. Introduction

Tabernaemontana longipes Donn. Sm. is one of about 110 species of *Tabernaemontana* in the Apocynaceae [1]. The plant ranges from Nicaragua to Colombia and Ecuador in very humid forests, secondary forests, as well as windy and cloud forests, 0-1900 m elevation. The inflorescences have many sweet-smelling flowers, which have white or cream-colored cylindrical corollas, 8-14 mm long, with brown lobes [2, 3]. Several alkaloids [4] and triterpenoids [5, 6] have been isolated and characterized from the plant. Apparently, however, the floral essential oil has not been previously examined.

2. Materials and methods

2.1 Plant Material

Flowers of *T. longipes* were collected on 9 May, 2009, from several treelets growing in the Monteverde region of northwestern Costa Rica (10° 18' 3.53" N, 84° 48' 39.43" W, 1350 m elevation). The plant was identified by William A. Haber. A voucher specimen has been deposited with the Missouri Botanical Garden herbarium (Haber Collection No. 7104). The fresh flowers (92.0 g) were hydrodistilled using a Likens-Nickerson apparatus with continuous extraction of the distillate with dichloromethane to give a clear colorless essential oil (15.4 mg).

2.2 Gas Chromatography – Mass Spectrometry

The floral essential oil of *T. longipes* was subjected to GC–MS analysis on an Agilent system consisting of a Model 6890 gas chromatograph, a Model 5973 mass selective detector (EIMS, electron energy = 70 eV, scan range = 45-400 amu, and scan rate = 3.99 scans/sec), and an Agilent ChemStation data system. The GC column was an HP-5ms fused silica capillary with a (5% phenyl)-methylpolysiloxane stationary phase (30 m \times 0.25 mm i.d., film thickness 0.25 μ m). The carrier gas was helium with a column head pressure of 48.7 kPa and flow rate of 1.0 ml/min. Inlet temperature was 200 °C and interface temperature was 280 °C. The GC oven temperature program was used as follows: 40 °C initial temperature held for 10 min; increased at 3 °C/min to 200 °C, then 2 °C/min to 280 °C. The sample was dissolved in CH₂Cl₂ to give a 1% w/v solution; 1 μ l injections using a splitless injection technique were used.

Identification of oil components was achieved based on their retention indices (RI, determined with reference to a homologous series of normal alkanes) and by comparison of their mass spectral fragmentation patterns with those reported in the literature [7] and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version C.00.01.08)]. The percentages of each component are reported as raw percentages without standardization.

Correspondence:

William N. Setzer
 Department of Chemistry,
 University of Alabama in
 Huntsville, Huntsville, AL 35899,
 USA
 Email: wsetzer@chemistry.uah.edu

3. Results & Discussion

A total of 37 compounds representing 86.6% of the essential oil composition were identified (Table 1). The major components in the floral essential oil were the monoterpene (*E*)- β -ocimene (30.0%), the sesquiterpene germacrene D (12.5%), and the normal alkane heneicosane (12.9%). Several members of the Apocynaceae have floral essential oils rich in (*E*)- β -ocimene, including *Funastrum odoratum*, *Gonolobus barbatus*, *Oxypetalum ostenii*,

Sarcostemma viminalis [8], *Fockea edulis*, *Hoya incrassata*, *Marsdenia linearis*, and *Secamone afzelii* [9]. Like *T. longipes*, *Catharanthus roseus* floral essential oil was rich in alkanes, particularly heneicosane and tricosane [10]. Germacrene D has apparently not been previously identified as an abundant component of Apocynaceae floral essential oils, but it is a major constituent of *Magnolia grandiflora* (Magnoliaceae) floral essential oil [11, 12].

Table 1: Chemical composition of *Tabernaemontana longipes* floral essential oil

RI	Compound	%
1001	Dehydro-1,8-cineole	1.9
1037	1,8-Cineole	0.1
1061	(<i>E</i>)- β -Ocimene	30.0
1293	Indole	2.2
1373	α -Copaene	0.9
1376	Daucene	trace
1382	β -Cubebene	0.4
1393	β -Elemene	2.2
1401	(<i>Z</i>)-Jasmone	0.3
1419	(<i>E</i>)-Caryophyllene	1.7
1437	α - <i>trans</i> -Bergamotene	2.2
1451	α -Humulene	0.3
1459	Sesquisabinene	0.4
1466	<i>cis</i> -Muurolo-4(14),5-diene	0.2
1486	Germacrene-D	12.5
1489	β -Selinene	1.3
1494	Ledene (= Viridiflorene)	0.7
1498	Bicyclogermacrene	1.3
1503	Isodaucene	0.4
1507	β -Bisabolene	1.1
1509	(<i>Z</i>)- α -Bisabolene	0.7
1520	δ -Cadinene	0.2
1560	(<i>E</i>)-Nerolidol	0.4
1573	Dendrolasin	2.0
1900	Nonadecane	0.6
2000	Eicosane	0.2
2100	Heneicosane	12.9
2115	Unidentified diterpenoid	2.5
2119	Unidentified diterpenoid	4.3
2200	Docosane	0.4
2300	Tricosane	3.5
2400	Tetracosane	0.1
2500	Pentacosane	1.2
2600	Hexacosane	0.1
2700	Heptacosane	2.6
2800	Octacosane	0.2
2832	Unidentified diterpenoid	5.7
2900	Nonacosane	1.3
2946	Unidentified diterpenoid	0.9
3000	Triacosane	trace
3100	Untriacontane	trace

Indole (2.2%) was the only nitrogen-containing compound identified in *T. longipes* floral essential oil. This compound has been identified in the floral volatiles of a few members of the

Apocynaceae, including *Apteranthes joannis* (11.2%), *Hoodia gordonii* (4.6%), *Orbea variegata* (5.7%) [13], *Oxypetalum ostenii* (1.5%), *Sarcostemma brevipedicellatum* (4.5%), *Sarcostemma*

viminale (14.3%)^[8], *Marsdenia engleriana* (3.0%)^[9], *Caralluma europaea* (0.8%)^[14], and *Plumeria rubra* (trace amount)^[15]. Interestingly, *T. longipes* floral essential oil was devoid of the “typical” essential oil components that contribute to floral fragrance, linalool and linalool derivatives, benzyl alcohol, and 2-phenylethanol, which have been observed in other floral essential oils from the Apocynaceae^[15-17]

4. Acknowledgments

I am grateful to the Monteverde Cloud Forest Preserve and the Tropical Science Center for permission to collect plant materials under a cooperative rights agreement and to the Commission for the Development of Biodiversity of Costa Rica’s Ministry of the Environment, Energy, and Telecommunications for Research Permit R-001-2006-OT-CONAGEBIO. I thank Dr. William A. Haber for plant identification and Dr. Bernhard Vogler for technical assistance with the GC-MS data collection.

5. References

- Mabberley DJ. *Mabberley’s Plant Book*. Edn 3, Cambridge University Press, UK, 2008, 837.
- Smith JD. Undescribed plants from Guatemala and other Central American Republics XIX. *Botanical Gazette* 1897; 24:389-398.
- Morales JF. Estudios en las Apocynaceae neotropicales XIX: La familia Apocynaceae s. str. (Apocynoideae, Rauvolfioideae) de Costa Rica. *Darwiniana* 2005; 43:90-191.
- Ciccio JF. Alcaloides de las semillas de *Tabernaemontana longipes* Donn. Smith. *Revista Latinoamericana de Quimica* 1979; 10:185.
- Ciccio JF, Hoet P. Algunos constituyentes de los frutos y de las hojas de *Tabernaemontana longipes* Donn Smith. *Revista Latinoamericana de Quimica* 1981; 12:88-90.
- Titant B, Germain G, Declercq JP, Meerssche VM, Ciccio JF, Hoet P. Crystal structure of baurenyl acetate from *Tabernaemontana longipes* Donn. Smith. *Bulletin des Sociétés Chimiques Belges* 1982; 91(2):117-121.
- Adams RP. *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*. Edn 4, Allured Publishing, Carol Stream, Illinois, USA, 2007.
- Jürgens A, Dötterl S, Liede-Schumann S, Meve U. Chemical diversity of floral volatiles in Asclepiadoideae-Asclepiadeae (Apocynaceae). *Biochemical Systematics and Ecology* 2008; 36:842-852.
- Jürgens A, Dötterl S, Liede-Schumann S, Meve U. Floral scent composition in early diverging taxa of Asclepiadoideae, and Secamonoideae (Apocynaceae). *South African Journal of Botany* 2010; 76:749-761.
- Pandey-Rai S, Mallavarapu GR, Naqvi AA, Yadav A, Rai SK, Srivastava S *et al.* Volatile components of leaves and flowers of periwinkle *Catharanthus roseus* (L.) G. Don from New Delhi. *Flavour and Fragrance Journal* 2006; 21:427-430.
- Davé PC, Vogler B, Setzer WN. Composition of the floral essential oil of *Magnolia grandiflora* L. (Magnoliaceae): Intraspecific and floral maturity variations. *Journal of Essential Oil-Bearing Plants* 2012; 15:694-702.
- Farag MA, Al-Mahdy DA. Comparative study of the chemical composition and biological activities of *Magnolia grandiflora* and *Magnolia virginiana* flower essential oils. *Natural Product Research* 2013; 27:1091-1097.
- Jürgens A, Dötterl S, Meve U. The chemical nature of fetid floral odours in stapeliads (Apocynaceae-Asclepiadoideae-Ceropegieae). *New Phytologist* 2006; 172:452-468.
- Formisano C, Senatore F, Porta DG, Scognamiglio M, Bruno M, Maggio A *et al.* Headspace volatile composition of the flowers of *Caralluma europaea* N.E.Br. (Apocynaceae). *Molecules* 2009; 14:4597-4613.
- Omata A, Yomogida K, Nakamura S, Hashimoto S, Arai T, Furukawa K. Volatile components of *Plumeria* flowers. Part 1. *Plumeria rubra* forma *acutifolia* (Poir.) Woodson cv. ‘Common Yellow’. *Flavour and Fragrance Journal* 1991; 6:277-279.
- Omata A, Nakamura S, Hashimoto S, Furukawa K. Volatile components of *Plumeria* flowers. Part 2. *Plumeria rubra* L. cv. ‘Irma Bryan’. *Flavour and Fragrance Journal* 1992; 7:33-35.
- Dung NX, Ngoc PH, Rang DD, Nhan NT, Klinkby N, Leclercq P. Chemical composition of the volatile concentrate from the flowers of Vietnamese *Alstonia scholaris* (L.) R.Br., Apocynaceae. *Journal of Essential Oil Research* 2001; 13:424-426.