



AkiNik

American Journal of Essential Oils and Natural Products

Available online at www.essencejournal.com

A
J
E
O
N
P

American
Journal of
Essential
Oils and
Natural
Products

ISSN: 2321-9114
AJEONP 2015; 3(2): 07-10
© 2015 AkiNik Publications
Received: 28-06-2015
Accepted: 30-07-2015

Prabodh Satyal

Department of Chemistry,
University of Alabama in Huntsville
Huntsville, AL 35899, USA.

William N. Setzer

Department of Chemistry,
University of Alabama in Huntsville
Huntsville, AL 35899, USA.

Chemical composition of *Cryptomeria japonica* leaf oil from Nepal

Prabodh Satyal, William N. Setzer

Abstract

Cryptomeria japonica is a commercially important timber tree throughout Asia, and the leaf essential oils from this tree have shown various biological activities. In this work, we have examined the chemical composition of the leaf essential oil of *C. japonica* from Nepal and compared the composition to those from other geographical locations. The leaf essential oil of *C. japonica* from Nepal was found to be qualitatively similar to leaf oils from other geographical locations, but very rich in both elemol and 16-kaurene, and may be a commercially-important source of these two phytochemicals.

Keywords: *Cryptomeria japonica*, chemical composition, leaf essential oil, elemol, kaurene, cluster analysis.

1. Introduction

Cryptomeria japonica (Thunb. ex L. f.) D. Don (Cupressaceae), known as “sugi” in Japan and “dhupi salla” in Nepal, is a commercially important tree in Japan, Korea, Taiwan, India, and China, and has been extensively cultivated worldwide. The wood of this tree, which has a fragrant odor and is insect resistant, has been used for construction, and the tree itself is valued in landscaping. There have been several recent reports on the biological activity of essential oils from *C. japonica*. The leaf oil from this plant has demonstrated insecticidal [1-3], antibacterial [4-6], antifungal [7, 8], cytotoxic [9], antiulcer [10], anxiolytic, and analgesic [11] activities. In this work, we have examined the chemical composition of the leaf essential oil of *C. japonica* from Nepal and compared the composition to those from other geographical locations.

2. Materials and Methods

2.1 Plant Material

The leaves of *C. japonica* were collected from the city of Kirtipur (27.67° N, 85.28° E, 1360 m above sea level) in Kathmandu district in Bagmati Zone in Nepal on 18 May 2011. The plant was identified by Nawal Shrestha, and a voucher specimen has been deposited in the herbarium of the Tribhuvan University Central Herbarium, Kirtipur, Nepal. The dry leaf sample (100 g) was crushed and hydrodistilled using a Clevenger type apparatus for 4 h to give 0.5 g of a clear pale yellow essential oil, which was stored at 4°C until analysis.

2.2 Gas Chromatographic – Mass Spectral Analysis

The essential oil from the leaves of *C. japonica* was analyzed by GC-MS using an Agilent 6890 GC with Agilent 5973 mass selective detector [MSD, operated in the EI mode (electron energy = 70 eV), scan range = 45-400 amu, and scan rate = 3.99 scans/sec], and an Agilent ChemStation data system. The GC column was an HP-5ms fused silica capillary with a (5% phenyl)-polymethylsiloxane stationary phase, film thickness of 0.25 µm, a length of 30 m, and an internal diameter of 0.25 mm. The carrier gas was helium with a column head pressure of 48.7 kPa and a flow rate of 1.0 mL/min. Injector temperature was 200 °C and detector temperature was 280 °C. The GC oven temperature program was used as follows: 40°C initial temperature, hold for 10 min; increased at 3 °C/min to 200 °C; increased 2°/min to 220 °C. A 1 % w/v solution of the sample in CH₂Cl₂ was prepared and 1 µL was injected using a splitless injection technique.

Identification of the oil components was based on their retention indices determined by reference to a homologous series of *n*-alkanes, and by comparison of their mass spectral fragmentation patterns with those reported in the literature [12] and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version

Correspondence:

Prabodh Satyal

Department of Chemistry,
University of Alabama in Huntsville
Huntsville, AL 35899, USA.

D.01.00)/Chem Station data system (G1701CA, version C.00.01.080)]. The percentages of each component are reported as raw percentages based on total ion current without standardization. The chemical composition of *C. japonica* leaf oil is summarized in Table 1.

2.3 Hierarchical Cluster Analysis

A total of 13 *C. japonica* leaf oil compositions from the published literature as well as the composition from this study were treated as operational taxonomic units (OTUs). The percentage composition of 25 essential oil components was

used to determine the chemical relationship between the various *C. japonica* essential oil samples by agglomerative hierarchical cluster (AHC) analysis using the XLSTAT software, version 2015.4.01. Pearson's correlation was selected as a measure of similarity, and the unweighted, pair-group method with arithmetic average (UPGMA) was used for cluster definition and to develop a dendrogram for the *C. japonica* samples. The compositions of the compounds used in the cluster analysis are listed in Table 2; the resulting dendrogram is shown in Figure 1.

Table 1: Chemical composition of *Cryptomeria japonica* leaf essential oil collected from Kirtipur, Nepal.

RI ^a	Compound	%	RI	Compound	%
936	α -Thujene	0.25	1515	γ -Cadinene	0.24
942	α -Pinene	4.20	1525	δ -Cadinene	1.45
954	Camphene	tr ^b	1557	Elemol	20.35
977	Sabinene	4.27	1620	10- <i>epi</i> - γ -Eudesmol	0.23
979	β -Pinene	0.14	1635	γ -Eudesmol	7.02
993	Myrcene	0.63	1644	τ -Cadinol	tr
1010	δ -3-Carene	0.10	1645	τ -Muurolool	tr
1017	α -Terpinene	0.51	1653	β -Eudesmol	4.99
1025	<i>p</i> -Cymene	0.33	1656	α -Eudesmol	4.74
1029	Limonene	1.13	1658	α -Cadinol	2.25
1059	γ -Terpinene	1.17	1670	Bulnesol	0.07
1088	Terpinolene	0.45	1677	Elemol acetate	0.14
1177	Terpinen-4-ol	0.91	1924	Unidentified diterpene ^c	1.17
1285	Bornyl acetate	0.12	1992	15-Kaurene	0.23
1393	β -Elemene	tr	2042	16-Kaurene	42.08
1478	γ -Muurolole	tr	2128	Nezukol	0.61
1501	α -Muurolole	0.21		Total Identified	98.83

^a RI = Retention index with respect to a homologous series of *n*-alkanes on an HP-5ms column.

^b tr = "trace" (< 0.05%)

^c MS: 258(9%), 257(66%), 187(11%), 175(13%), 161(19%), 137(37%), 133(32%), 123(40%), 121(35%), 119(42%), 109(54%), 107(49%), 105(49%), 95(93%), 93(97%), 91(86%), 81(100%), 79(89%), 69(86%), 67(76%), 55(73%), 53(35%).

3. Results and Discussion

The leaf oil of *C. japonica* from Nepal was dominated by the diterpene 16-kaurene (42.1%), the sesquiterpenoids elemol (20.3%), γ -eudesmol (7.0%), β -eudesmol (5.0%), and α -eudesmol (4.7%), and the monoterpenes α -pinene (4.2%) and sabinene (4.3%). The chemical composition of the Nepalese leaf oil is qualitatively similar to leaf oils from other geographical locations (see Table 2). Thus, *C. japonica* leaf oil is generally characterized as having large concentrations of 16-kaurene, elemol, and isomers of eudesmol, with lesser

amounts of sabinene, α -pinene, and terpinene-4-ol. The cluster analysis (Figure 1) reveals the similarities in most of the leaf oils (chemotype A) with the exception of one sample from Korea [4], which had only a small concentration of 16-kaurene (1.3%), no γ -eudesmol, but a relatively large concentration of α -terpineol (6.1%) (chemotype B), and samples from Reunion Island [13] and Japan [14], which were rich in α -pinene and sabinene, but poor in γ -eudesmol (chemotype C)

Table 2: *Cryptomeria japonica* leaf oil compositions used for the cluster analysis.

Compound	Nepal ^a	Korea 1 [5]	Korea 2 [4]	Korea 3 [6]	Korea 4 [8]	Taiwan 1 [15]	Taiwan 2 [11]	Taiwan 3 [2]	Taiwan 4 [2]	Taiwan 5 [2]	Taiwan 6 [3]	China [16]	Reunion Island [13]	Japan [14]
α -Thujene	0.25	0.00	1.32	0.54	0.57	0.00	0.79	1.21	1.28	0.92	0.46	0.47	0.00	0.00
α -Pinene	4.20	3.48	6.07	4.21	2.96	6.53	5.62	4.77	4.92	4.40	8.54	8.00	37.20	30.72
Camphene	0.03	0.50	0.42	0.97	0.61	0.63	0.96	0.46	0.46	0.46	3.36	0.81	5.38	3.72
Sabinene	4.27	8.86	8.86	11.06	5.06	10.29	9.37	10.21	10.83	6.81	3.80	0.00	18.80	11.30
β -Pinene	0.14	0.00	0.46	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.00	0.66	1.57	1.80
Myrcene	0.63	1.17	0.00	2.26	1.26	1.71	2.59	1.14	1.16	1.03	1.55	0.00	6.53	9.20
δ -3-Carene	0.10	1.73	0.42	1.61	0.62	13.05	9.66	1.24	1.03	0.77	1.50	0.53	5.60	5.42
α -Terpinene	0.51	0.97	2.58	0.90	1.42	2.46	1.90	0.77	1.19	0.86	0.91	0.05	4.50	6.28
<i>p</i> -Cymene	0.33	0.00	1.14	0.00	0.00	0.52	0.21	2.31	1.88	2.65	0.00	0.12	2.90	1.62
Limonene	1.13	1.22	3.43	3.55	1.89	3.67	5.26	2.09	2.03	2.35	6.81	1.63	4.84	12.20
γ -Terpinene	1.17	1.92	4.71	1.78	2.15	4.25	3.08	1.82	2.35	2.01	1.36	0.63	7.73	11.30
Terpinolene	0.45	1.13	0.00	0.88	0.93	1.85	1.58	0.82	1.00	0.84	0.62	0.28	2.30	3.72
Terpinen-4-ol	0.91	4.10	9.77	1.68	4.63	5.99	9.06	6.19	6.50	8.34	2.01	0.69	5.86	17.57
α -Terpineol	0.00	0.00	6.13	0.00	0.29	0.00	0.51	0.00	0.07	0.18	0.23	0.00	0.47	1.20
Bornyl acetate	0.12	0.00	3.14	0.00	0.71	0.70	0.76	0.89	1.11	1.34	3.84	0.51	1.60	1.85
β -Elemene	0.03	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.00	0.00	0.00	5.87	0.14	0.00
α -Muurolole	0.21	0.52	0.00	0.47	0.28	0.00	0.00	0.24	0.22	0.18	0.00	0.33	0.15	0.24
γ -Cadinene	0.24	0.00	0.00	0.00	0.21	0.00	0.00	0.22	0.20	0.15	0.00	1.47	0.25	0.94

δ-Cadinene	1.45	3.31	2.29	1.61	1.78	0.00	0.53	1.14	1.05	0.84	0.41	1.73	0.25	0.94
Elemol	20.35	10.88	11.17	8.60	6.87	13.93	18.22	19.05	18.41	18.30	18.29	20.12	11.75	4.25
γ-Eudesmol	7.02	9.41	0.00	2.88	19.02	0.00	0.00	6.27	6.40	7.16	8.23	4.12	0.05	0.00
β-Eudesmol	4.99	5.13	3.90	4.82	5.98	4.52	0.00	5.90	5.77	5.92	4.84	5.03	3.40	1.66
α-Eudesmol	4.74	5.26	2.58	3.53	7.86	0.00	5.70	5.90	5.77	5.92	6.50	5.62	2.85	1.65
α-Cadinol	2.25	2.20	4.28	1.48	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.39	0.68
16-Kaurene	42.08	17.20	1.27	31.45	26.31	21.74	11.63	20.44	19.53	20.69	23.29	14.84	10.00	4.80

^a This work




Fig 1: Dendrogram obtained from agglomerative hierarchical cluster of 14 *Cryptomeria japonica* essential oil compositions

4. Conclusions

The leaf essential oil of *Cryptomeria japonica* from Kirtipur, Nepal was found to be rich in both elemol and 16-kaurene, and may be a commercially-important source of these two phytochemicals.

References

1. Wang SY, Lai WC, Chu FH, Lin CT, Shen SY, Chang ST. Essential oil from the leaves of *Cryptomeria japonica* acts as a silverfish (*Lepisma saccharina*) repellent and insecticide. *Journal of Wood Science*, 2006; 52: 522-526.
2. Cheng SS, Chua MT, Chang EH, Huang CG, Chen WJ, Chang ST. Variations in insecticidal activity and chemical compositions of leaf essential oils from *Cryptomeria japonica* at different ages. *Bioresource Technology*, 2009; 100:465-470.
3. Cheng SS, Lin CY, Chung MJ, Chang ST. Chemical composition and antitermitic activity against *Coptotermes formosanus* Shiraki of *Cryptomeria japonica* leaf essential

- oil. *Chemistry & Biodiversity*, 2012; 9:352-358.
4. Cha JD, Jeong MR, Jeong SI, Moon SE, Kil BS, Yun SI, Lee KY, Song YH. Chemical composition and antimicrobial activity of the essential oil of *Cryptomeria japonica*. *Phytotherapy Research*, 2007; 21:295-299.
5. Yoon WJ, Kim SS, Oh TH, Lee NH, Hyun CG. *Cryptomeria japonica* essential oil inhibits the growth of drug-resistant skin pathogens and LPS-induced nitric oxide and pro-inflammatory cytokine production. *Polish Journal of Microbiology*, 2009; 58: 61-68.
6. Lee JH, Lee BK, Kim JH, Lee SH, Hong SK. Comparison of chemical compositions and antimicrobial activities of essential oils from three conifer trees; *Pinus densiflora*, *Cryptomeria japonica*, and *Chamaecyparis obtusa*. *Journal of Microbiology and Biotechnology*, 2009; 19: 391-396.
7. Cheng SS, Lin HY, Chang ST. Chemical composition and antifungal activity of essential oils from different tissues of Japanese cedar (*Cryptomeria japonica*). *Journal*

- of Agricultural and Food Chemistry, 2005; 53:614-619.
8. Oh HJ, Ahn HM, So KH, Kim SS, Yun PY, Jeon GL *et al.* Chemical and antimicrobial properties of essential oils from three coniferous trees *Abies koreana*, *Cryptomeria japonica*, and *Torreya nucifera*. *Journal of Applied Biological Chemistry*, 2007; 50:164-169.
 9. Cha JD, Kim JY. Essential oil from *Cryptomeria japonica* induces apoptosis in human oral epidermoid carcinoma cells via mitochondrial stress and activation of caspases. *Molecules*, 2012; 17:3890-3901.
 10. Matsunaga T, Hasegawa C, Kawasuji T, Suzuki H, Saito H, Sagioka T, Takahashi R, Tsukamoto H, Morikawa T, Akiyama T. Isolation of the antiulcer compound in essential oil from the leaves of *Cryptomeria japonica*. *Biological and Pharmaceutical Bulletin*, 2000; 23:595-598.
 11. Cheng WW, Liu CT, Chu FH, Chang ST, Wang SY. Neuropharmacological activities of phytoncide released from *Cryptomeria japonica*. *Journal of Wood Science*, 2009; 55:27-31.
 12. Adams RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, 4th Edition. Allured Publishing, Carol Stream, Illinois, USA, 2007.
 13. Vernin G, Metzger J, Mondon JP, Pieribattesti JC. GC/MS analysis of the leaf oil of *Cryptomeria japonica* D. Don from Reunion Island. *Journal of Essential Oil Research*, 1991; 3:197-207.
 14. Yatagai M, Sato T. Terpenes of leaf oils from conifers. *Biochemical Systematics and Ecology*, 1986; 14:469-478.
 15. Gu HJ, Cheng SS, Lin CY, Huang CG, Chen WJ, Chang ST. Repellency of essential oils of *Cryptomeria japonica* (Pinaceae) against adults of the mosquitoes *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Journal of Agricultural and Food Chemistry*, 2009; 57:11127-11133.
 16. Xie Y, Wang J, Yang F, Lei C. Comparative analysis of essential oil components of two *Cryptomeria* species from China. *Industrial Crops and Products*, 2011; 34:1226-1230.