



AkiNik

# American Journal of Essential Oils and Natural Products

Available online at [www.essencejournal.com](http://www.essencejournal.com)

A  
J  
E  
O  
N  
P  
American  
Journal of  
Essential  
Oils and  
Natural  
Products

ISSN: 2321-9114  
 AJEONP 2017; 5(4): 26-29  
 © 2017 AkiNik Publications  
 Received: 19-08-2017  
 Accepted: 20-09-2017

**Jonathan D Craft**  
 Department of Chemistry,  
 University of Alabama in  
 Huntsville Huntsville, AL 35899,  
 USA

**William N Setzer**  
 (A). Department of Chemistry,  
 University of Alabama in  
 Huntsville Huntsville, AL 35899,  
 USA  
 (B). Aromatic Plant Research  
 Center, 615 St. George Square  
 Court, Suite 300  
 Winston-Salem, NC 27103, USA

**Correspondence**  
**William N Setzer**  
 (A). Department of Chemistry,  
 University of Alabama in  
 Huntsville Huntsville, AL 35899,  
 USA  
 (B). Aromatic Plant Research  
 Center, 615 St. George Square  
 Court, Suite 300  
 Winston-Salem, NC 27103, USA

## Leaf essential oil composition of *Tsuga canadensis* growing wild in North Alabama and Northwest Georgia

**Jonathan D Craft and William N Setzer**

### Abstract

*Tsuga canadensis* was an important traditional medicine for Native Americans for the treatment of rheumatism. Leaves of *T. canadensis* were collected from two disjunctive populations: one from north Alabama and the other from northwest Georgia. The leaf essential oils were obtained by hydrodistillation and analyzed by GC-FID and GC-MS. The major components in the leaf oils were isobornyl acetate,  $\alpha$ -pinene, camphene, limonene, piperitone, and manool. The presence of these constituents, which had previously shown analgesic and anti-inflammatory activities, support the traditional uses of this plant to treat rheumatism.

**Keywords:** Native American traditional medicine, isobornyl acetate,  $\alpha$ -pinene, camphene, limonene, piperitone, manool

### 1. Introduction

The genus *Tsuga* (Pinaceae) is comprised of nine species, four of which occur in North America [1]. *Tsuga canadensis* (L.) Carrière, “eastern hemlock”, was an important traditional medicine for Native Americans. The Abenaki and the Iriquois people consumed a decoction of the leaves to treat rheumatism; a decoction of the twigs was used by the Delaware Native Americans as an herbal steam to treat rheumatism [2]. In 1951, Shaw obtained the steam-distilled essential oil of *T. canadensis*, and was able to isolate and identify seven monoterpene hydrocarbons: Tricyclene,  $\alpha$ -pinene, camphene,  $\beta$ -pinene, myrcene,  $\alpha$ -phellandrene, and limonene [3]. More recently, there have been several headspace solid-phase microextractions [4-7] and solvent extractions [8] along with mass spectral analyses of the leaves of *T. canadensis*. In addition to the seven major monoterpene hydrocarbons, the oxygenated monoterpenoids piperitone and isobornyl acetate, and the sesquiterpene hydrocarbons  $\beta$ -caryophyllene and  $\alpha$ -humulene, were also found to be major volatile components. In this work, we report the chemical compositions of the leaf essential oils of two different specimens of *T. canadensis* collected in northern Alabama and northwestern Georgia.

### 2. Materials and Methods

#### 2.1 Plant Material

Branches of *T. canadensis* were collected from north Alabama (34.308° N, 87.502° E, 230 m elevation) on November 1, 2017, and from northwestern Georgia (34.819° N, 85.495° E, 546 m elevation) on November 6, 2017. The fresh leaves were removed from the branches, stored at -20 °C overnight, and hydrodistilled using a Likens-Nickerson apparatus with continuous extraction with dichloromethane for four hours.

#### 2.2 Gas Chromatography and Gas Chromatography-Mass Spectrometry

The leaf essential oils of *T. canadensis* were analyzed by gas chromatography – flame ionization detection (GC-FID, Agilent 6890 GC with Agilent FID) and gas chromatography – mass spectrometry (GC-MS, Agilent 6890 GC, Agilent 5973 MSD, HP-5ms capillary column) as described previously [9].

### 3. Results and Discussion

The masses of the frozen leaves prior to extraction were 36.1 g and 33.0 g for the Alabama and Georgia samples, respectively. Hydrodistillation yielded 149 mg (Alabama) and 136 mg

(Georgia) of essential oils. The leaf essential oil compositions of *T. canadensis* from Alabama and Georgia are compiled in Table 1. The oils were dominated by monoterpene hydrocarbons (44.6 and 27.4%, respectively) and oxygenated monoterpenoids (48.4 and 51.9%, respectively). The major components of *T. canadensis* leaf essential oils were tricyclene (4.5% in the oil from Alabama),  $\alpha$ -pinene (13.9 and

5.4% in the oils from Alabama and Georgia, respectively), camphene (13.3 and 3.4%, respectively), limonene (6.0 and 7.0%, respectively), piperitone (4.3 and 7.7%, respectively), bornyl acetate (38.6 and 37.0%, respectively),  $\beta$ -caryophyllene (3.4% in the oil from Georgia),  $\alpha$ -humulene (4.4% in the oil from Georgia), and the diterpenoid manool (6.8% in the oil from Georgia).

**Table 1:** Chemical compositions of the leaf essential oils of *Tsuga canadensis* collected from Bankhead National Forest, Alabama, and Rising Fawn, Georgia.

RI	Compound	Percent Composition	
		Alabama	Georgia
851	(2E)-Hexenal	0.6	0.2
881	Santene	tr	---
924	Tricyclene	4.5	0.6
929	$\alpha$ -Thujene	0.2	0.1
936	$\alpha$ -Pinene	13.9	5.4
948	Camphene	13.3	3.4
952	Thuja-2,4(10)-diene	tr	tr
970	Sabinene	0.1	0.1
973	$\beta$ -Pinene	2.0	2.2
989	Myrcene	1.6	1.7
1001	$\alpha$ -Phellandrene	0.7	3.7
1006	$\delta$ -3-Carene	0.1	0.1
1013	$\alpha$ -Terpinene	0.3	0.3
1021	<i>p</i> -Cymene	0.2	0.7
1027	Limonene	6.0	7.0
1028	1,8-Cineole	0.4	---
1038	(Z)- $\beta$ -Ocimene	tr	tr
1049	(E)- $\beta$ -Ocimene	tr	tr
1058	$\gamma$ -Terpinene	0.5	0.6
1066	<i>cis</i> -Sabinene hydrate	tr	0.1
1089	Terpinolene	1.3	1.5
1098	<i>trans</i> -Sabinene hydrate	tr	0.1
1103	Linalool	tr	0.1
1114	<i>endo</i> -Fenchol	tr	tr
1123	<i>cis-p</i> -Menth-2-en-1-ol	0.2	0.7
1128	$\alpha$ -Campholenal	tr	tr
1139	<i>trans</i> -Pinocarveol	tr	tr
1142	<i>trans-p</i> -Menth-2-en-1-ol	0.1	0.5
1145	Camphor	0.6	0.6
1147	<i>trans</i> -Verbenol	tr	0.1
1149	Camphene hydrate	0.3	0.1
1151	$\alpha$ -Phellandrene-8-ol	tr	tr
1158	Isoborneol	tr	---
1164	Pinocarvone	tr	tr
1167	Borneol	0.7	0.9
1170	<i>p</i> -Mentha-1,5-dien-8-ol	0.1	0.1
1175	<i>cis</i> -Pinocamphone	tr	tr
1180	Terpinen-4-ol	0.8	1.5
1194	$\alpha$ -Terpineol	1.2	0.8
1198	<i>cis</i> -Piperitol	0.1	0.2
1210	Verbenone	0.1	0.2
1211	<i>trans</i> -Piperitol	0.1	0.2
1234	Citronellol	tr	0.1
1238	Thymol methyl ether	0.1	---
1253	Piperitone	4.3	7.7
1286	Isobornyl acetate	38.6	37.0
1290	Thymol	0.5	0.4
1295	<i>trans</i> -Pinocarvyl acetate	tr	0.1
1327	Myrtenyl acetate	0.1	0.2
1350	$\alpha$ -Cubebene	tr	0.1
1358	Citronellyl acetate	0.1	tr
1369	Neryl acetate	tr	---
1370	$\alpha$ -Ylangene	tr	tr
1375	$\alpha$ -Copaene	tr	0.1
1383	$\beta$ -Bourbonene	tr	tr
1384	<i>trans</i> -Myrtanyl acetate	tr	0.1

1392	7- <i>epi</i> -Sesquithujene	---	tr
1392	$\beta$ -Cubebene	---	tr
1418	$\beta$ -Caryophyllene	1.0	3.4
1427	$\beta$ -Copaene	tr	tr
1437	$\alpha$ - <i>trans</i> -Bergamotene	tr	0.1
1452	$\alpha$ -Humulene	1.6	4.4
1460	Sesquisabinene	tr	tr
1462	<i>cis</i> -Cadina-1(6),4-diene	tr	tr
1473	<i>trans</i> -Cadina-1(6),4-diene	tr	tr
1476	$\gamma$ -Muuroleone	0.2	0.4
1479	Germacrene D	0.1	0.4
1484	$\beta$ -Selinene	0.1	0.3
1490	<i>trans</i> -Muurolo-4(14),5-diene	tr	tr
1493	$\alpha$ -Selinene	tr	0.1
1494	$\gamma$ -Amorphene	tr	0.1
1501	$\alpha$ -Muuroleone	0.1	0.1
1507	Epizonarene	tr	---
1511	$\beta$ -Bisabolene	tr	0.2
1513	$\gamma$ -Cadinene	0.1	0.5
1517	( <i>Z</i> )- $\gamma$ -Bisabolene	tr	0.3
1524	$\delta$ -Cadinene	0.3	0.8
1534	( <i>E</i> )- $\gamma$ -Bisabolene	tr	0.3
1542	$\alpha$ -Calacorene	tr	tr
1567	( <i>E</i> )-Nerolidol	0.5	0.8
1581	Caryophyllene oxide	0.2	0.2
1607	Humulene epoxide II	0.2	0.2
1642	$\tau$ -Cadinol	0.2	0.3
1647	$\alpha$ -Muurolol	0.1	0.1
1654	$\alpha$ -Cadinol	0.2	0.2
1688	$\alpha$ -Bisabolol	tr	0.1
1726	(2 <i>Z</i> ,6 <i>E</i> )-Farnesol	tr	0.1
2053	Manool	1.6	6.8
	Monoterpene hydrocarbons	44.6	27.4
	Oxygenated monoterpenoids	48.4	51.9
	Sesquiterpene hydrocarbons	3.4	11.5
	Oxygenated sesquiterpenoids	1.3	2.2
	Diterpenoids	1.6	6.8
	Others	0.6	0.2
	Total Identified	99.9	99.6

The compositions of *T. canadensis* essential oils are consistent with the headspace solid-phase microextractions reported earlier [4-7]: Tricyclene (3.1-7.8%),  $\alpha$ -pinene (10.4-23.7%), camphene (7.8-15.9%), limonene (0.7-6.0%), piperitone (up to 5.9%), isobornyl acetate (26.8-50.7%),  $\beta$ -caryophyllene (1.3-6.0%), and  $\alpha$ -humulene (up to 9.2%). However, these headspace techniques did not reveal less volatile constituents such as oxygenated sesquiterpenoids and the diterpenoid manool, which were observed in the essential oils in this study.

Several monoterpenes have shown analgesic [10, 11] and anti-inflammatory [12-14] effects, including  $\alpha$ -pinene, myrcene, and limonene. In addition, bornyl acetate [15-17] has shown anti-inflammatory activity,  $\beta$ -caryophyllene has demonstrated analgesic [18-21] and anti-inflammatory [14, 22] activities,  $\alpha$ -humulene has shown anti-inflammatory effects [23], and manool was also shown to be anti-inflammatory [24]. Thus, the major components of *T. canadensis* essential oil show analgesic and anti-inflammatory activities, which are consistent with the traditional uses of this plant to treat rheumatism.

#### Acknowledgments

This work was carried out as part of the activities of the Aromatic Plant Research Center (APRC, <https://aromaticplant.org/>). WNS is grateful to dōTERRA International (<https://www.doterra.com/US/en>) for financial

support of the APRC.

#### Conflicts of Interest

The authors declare no conflicts of interest. The funding sponsor, dōTERRA International, played no role in the design of the study; in the collection, analysis, or interpretation of the data; conclusions of the study; or in the decision to publish the results.

#### References

- Mabberley DJ. *Mabberley's Plant-Book*, 3rd ed. Cambridge University Press, Cambridge, UK, 2008.
- Moerman DE. *Native American Ethnobotany*. Timber Press, Portland, OR, USA, 1998.
- Shaw AC. The essential oil of *Tsuga canadensis* (L.) Carr. *Journal of the American Chemical Society*, 1951; 73:2859-2861.
- Lagalante AF, Montgomery ME. Analysis of terpenoids from hemlock (*Tsuga*) species by solid-phase microextraction / gas chromatography / ion-trap mass spectrometry. *Journal of Agricultural and Food Chemistry*. 2003; 51:2115-2120.
- Lagalante AF, Lewis N, Montgomery ME, Shields KS. Temporal and spatial variation of terpenoids in eastern hemlock (*Tsuga canadensis*) in relation to feeding by *Adelgestsugae*. *Journal of Chemical Ecology*. 2006; 32:2389-2403.

6. Lagalante AF, Montgomery ME, Calvosa FC, Mirzabeigi MN. Characterization of terpenoid volatiles from cultivars of eastern hemlock (*Tsuga canadensis*). Journal of Agricultural and Food Chemistry. 2007; 55:10850-10856.
7. Kılıç Ö, Kocak A. Volatile constituents of *Juniperus communis* L., *Taxus canadensis* Marshall. and *Tsuga canadensis* (L.) Carr. from Canada. Journal of Agricultural Science and Technology B. 2014; 4:135-140.
8. McKenzie EA, Elkinton JS, Casagrande RA, Preisser EI, Mayer M. Terpene chemistry of eastern hemlocks resistant to hemlock woolly adelgid. Journal of Chemical Ecology. 2014; 40:1003-1012.
9. Satyal P, Craft JD, Dosoky NS, Setzer WN. The chemical compositions of the volatile oils of garlic (*Allium sativum*) and wild garlic (*Allium vineale*). Foods. 2017; 6:63.
10. de Sousa DP. Analgesic-like activity of essential oils constituents. Molecules. 2011; 16:2233-2252.
11. Guimarães AG, Quintans JSS, Quintans-Júnior LJ. Monoterpenes with analgesic activity-A systematic review. Phytherapy Research. 2013; 27:1-15.
12. Salminen A, Lehtonen M, Suuronen T, Kaarniranta K, Huuskonen J. Terpenoids: Natural inhibitors of NF-κB signaling with anti-inflammatory and anticancer potential. Cellular and Molecular Life Sciences. 2008; 65:2979-2999.
13. da Silveira e Sá RdC, Andrade LN, de Sousa DP. A review on anti-inflammatory activity of monoterpenes. Molecules. 2013; 18:1227-1254.
14. Rufino AT, Ribeiro M, Sousa C, Judas F, Salgueiro L, Cavaleiro C *et al.* Evaluation of the anti-inflammatory, anti-catabolic and pro-anabolic effects of *E*-caryophyllene, myrcene and limonene in a cell model of osteoarthritis. European Journal of Pharmacology. 2015; 750:141-150.
15. Wu X, Li X, Xiao F, Zhang Z, Xu Z, Wang H. [Studies on the analgesic and anti-inflammatory effect of bornyl acetate in volatile oil from *Amomum villosum*]. Zhong Yao Cai. 2004; 27:438-439.
16. Chen N, Sun G, Yuan X, Hou J, Wu Q, Soromou LW *et al.* Inhibition of lung inflammatory responses by bornyl acetate is correlated with regulation of myeloperoxidase activity. Journal of Surgical Research. 2014; 186:436-445.
17. Yang H, Zhao R, Chen H, Jia P, Bao L, Tang H. Bornyl acetate has an anti-inflammatory effect in human chondrocytes via induction of IL-11. International Union of Biochemistry and Molecular Biology. 2014; 67:854-859.
18. Ghelardini C, Galeotti N, Mannelli LDC, Mazzanti G, Bartolini A. Local anaesthetic activity of β-caryophyllene. Il Farmaco. 2001; 56:387-389.
19. Gertsch J, Leonti M, Raduner S, Racz I, Chen JZ, Xie XQ *et al.* Beta-caryophyllene is a dietary cannabinoid. Proceedings of the National Academy of Sciences of the United States of America. 2008; 105:9099-9104.
20. Bakır B, Him A, Özbek H, Düz E, Tütüncü M. Investigation of the anti-inflammatory and analgesic activities of β-caryophyllene. International Journal of Essential Oil Therapeutics. 2008; 2:41-44.
21. Paula-Freire LIG, Andersen ML, Gama VS, Molska GR, Carlini ELA. The oral administration of *trans*-caryophyllene attenuates acute and chronic pain in mice. Phytomedicine. 2014; 21:356-362.
22. Klauke A, Racz I, Pradier B, Markert A, Zimmer AM, Gertsch J *et al.* The cannabinoid CB2 receptor-selective phytocannabinoid beta-caryophyllene exerts analgesic effects in mouse models of inflammatory and neuropathic pain. European Neuropsychopharmacology. 2014; 24:608-620.
23. Fernandes ES, Passos GF, Medeiros R, da Cunha FM, Ferreira J, Campos MM *et al.* Anti-inflammatory effects of compounds alpha-humulene and (-)-*trans*-caryophyllene isolated from the essential oil of *Cordia verbenacea*. European Journal of Pharmacology. 2007; 569:228-236.
24. Li YL, Yang XW, LiSM, Shen YH, Zeng HW, Liu XH *et al.* Terpenoid constituents of *Abies chensiensis* with potential anti-inflammatory activity. Journal of Natural Products. 2009; 72:1065-1068.