



AkiNik

American Journal of Essential Oils and Natural Products

Available online at www.essencejournal.com

A
J
E
O
N
P
American
Journal of
Essential
Oils and
Natural
Products

ISSN: 2321-9114
AJEONP 2018; 6(3): 27-30
© 2018 AkiNik Publications
Received: 28-05-2018
Accepted: 29-06-2018

Andrea Vargas Suarez
Oeste de la Fabrica de Quesos de
Monteverde, Monteverde,
Puntarenas, 60109 Costa Rica

Prabodh Satyal
Aromatic Plant Research Center,
George Square Court, Suite,
Winston-Salem 27103, USA

William N Setzer
1) Aromatic Plant Research
Center, George Square Court,
Suite, Winston-Salem 35899,
USA
2) Department of Chemistry,
University of Alabama in
Huntsville Huntsville, AL 35899,
USA

Correspondence
William N Setzer

1) Aromatic Plant Research
Center, George Square Court,
Suite, Winston-Salem 35899,
USA
2) Department of Chemistry,
University of Alabama in
Huntsville Huntsville, AL 35899,
USA

Volatile Components of the Wood of Spanish Cedar, *Cedrela odorata*, from Costa Rica

Andrea Vargas Suarez, Prabodh Satyal, William N Setzer

Abstract

Cedrela odorata, “cedro amargo”, is a valuable lumber tree, which has fragrant smelling wood. The essential oil from the wood of *Cedrela odorata* from Guanacaste, Costa Rica, was obtained by hydrodistillation and analyzed by gas chromatography – mass spectrometry. The wood oil was rich in sesquiterpene hydrocarbons, including δ -cadinene (26%), β -curcumene (13%), and calarene (6%). In comparison, a commercial *C. odorata* wood oil sample from Colombia was dominated by δ -cadinene (53%), along with germacrene D (4%). The pleasant aroma and chemical components of *C. odorata* wood essential oil suggest that it may be an important resource for the cosmetic and flavor industries.

Keywords: Essential oil composition; δ -cadinene; β -curcumene; calarene

1. Introduction

Cedrela odorata L., “cedro amargo” (Meliaceae), is a large canopy tree that ranges from Mexico, through Central America and the West Indies, and into South America, including Peru, Brazil, and northern Argentina [1-3]. The wood is soft and light, easily worked and in much demand for carpentry, musical instruments, floors, ceilings, overlays, and crafts [3], resulting in overexploitation over much of its natural range [2]. The heartwood of *C. odorata* contains an aromatic, antifungal, and insect-repelling resin [4]. There have been previous reports on essential oils of *C. odorata* from the leaves from Nigeria [5] and Brazil [6], the bark from São Tomé y Príncipe [7] and Costa Rica [8], the stems from Brazil [6], and commercial wood oil [9]. In this work, we present the composition of the wood essential oil from *C. odorata* from Guanacaste Province, Costa Rica, and a commercial wood essential oil sample from Colombia.

2. Materials and Methods

2.1 Essential Oils

Wood shavings of *C. odorata* were obtained from the workshop of Leonardo Vargas Garcia, La Cruz, Abangares, Guanacaste, Costa Rica. Two different samples of 500 g of wood were hydrodistilled using a Mountain Home Biological SL-SS 20 L stainless steel essential oil distiller to give the wood oils in 1% yield. The wood essential oil from Colombia was a commercial sample provided by Daniel Sheffield (Essential Oil Quality Alliance, Charlestown, Indiana, USA).

2.2 Gas Chromatography – Mass Spectrometry

The *C. odorata* wood essential oils were analyzed by gas chromatography-mass spectrometry (GC-MS) using a Shimadzu GCMS-QP2010 Ultra operated in the electron impact (EI) mode (electron energy = 70 eV), scan range = 40–400 atomic mass units, scan rate = 3.0 scans/s, and GC-MS solution software v. 4.20 (Shimadzu Scientific Instruments, Columbia, MD, USA). The GC column was a ZB-5 fused silica capillary column (Phenomenex, Torrance, CA, USA) with a (5% phenyl)-polymethylsiloxane stationary phase and a film thickness of 0.25 μ m. The carrier gas was helium with a column head pressure of 552 kPa and flow rate of 1.37 mL/min. The injector temperature was 260 °C and the ion source temperature was 260 °C. The GC oven temperature program was programmed for 50 °C initial temperature, temperature increased at a rate of 2 °C/min to 260 °C. A 5% w/v solution of the sample in CH₂Cl₂ was prepared and 0.1 μ L was injected with a splitting mode (30:1). Identification of the oil components was based on their retention indices determined by reference to a homologous series of *n*-alkanes, and by comparison of their mass spectral fragmentation patterns with those reported in the literature

[10]. and stored in our in-house library [11].

3. Results and Discussion

The yellow, fragrant-smelling wood essential oils of *C. odorata* were obtained in 1% yield. The chemical compositions of the *C. odorata* wood essential oils from Costa Rica and from Colombia are compiled in Table 1. The *C. odorata* wood essential oils from Costa Rica were rich in sesquiterpene hydrocarbons δ -cadinene (26%), β -curcumene

(13%), and calarene (6%), as well as the sesquiterpene alcohol α -cadinol (5%). In contrast, the wood oil from Colombia was dominated by δ -cadinene (53%), along with germacrene D (4%). Unfortunately, the chemical composition percentages were not reported in a previously published *C. odorata* wood oil [9]. From the published chromatogram, however, the major components were δ -cadinene, α -copaene, and α -cubebene. The wood oil samples in this current work showed very little α -copaene or α -cubebene.

Table 1: Chemical compositions (%) of the wood essential oils of *Cedrela odorata*.

RI ^a	Compound	Costa Rica #1	Costa Rica #2	Colombia
931	α -Pinene	0.1	0.1	0.2
948	Camphene	tr ^b	tr	---
971	Sabinene	tr	tr	---
977	β -Pinene	tr	tr	---
987	Myrcene	tr	tr	tr
1006	α -Phellandrene	tr	tr	---
1024	<i>p</i> -Cymene	tr	tr	tr
1028	Limonene	tr	tr	3.2
1030	β -Phellandrene	tr	tr	---
1031	1,8-Cineole	tr	tr	0.4
1044	(<i>E</i>)- β -Ocimene	tr	tr	---
1057	γ -Terpinene	tr	tr	---
1084	Terpinolene	tr	tr	---
1099	Linalool	tr	tr	---
1152	Citronellal	tr	tr	---
1298	Carvacrol	tr	tr	---
1330	Bicycloelemene	tr	tr	---
1333	δ -Elemene	tr	tr	0.5
1345	α -Cubebene	tr	tr	0.3
1367	α -Ylangene	---	---	0.6
1374	α -Copaene	0.3	0.2	1.3
1386	β -Cubebene	---	---	0.3
1388	β -Elemene	0.6	0.5	1.1
1405	α -Gurjunene	0.1	0.1	---
1413	<i>cis</i> - α -Bergamotene	2.7	2.6	0.9
1415	2- <i>epi</i> - β -Furberene	0.2	0.2	---
1417	β -Caryophyllene	---	---	0.1
1418	Aristolene	1.9	1.8	---
1424	β -Cedrene	0.8	0.7	---
1427	γ -Elemene	---	---	0.7
1428	Aromadendrene	0.4	0.4	---
1431	<i>trans</i> - α -Bergamotene	---	---	tr
1432	β -Gurjunene (Calarene)	5.5	5.7	---
1433	α -Guaiene	---	---	tr
1435	α -Maaliene	3.4	3.4	---
1439	6,9-Guaiadiene	---	---	1.7
1439	(<i>Z</i>)- β -Farnesene	0.4	0.3	---
1444	<i>cis</i> -Muurolo-3,5-diene	0.1	0.1	---
1448	<i>trans</i> -Muurolo-3,5-diene	0.2	0.2	0.5
1453	(<i>E</i>)- β -Farnesene	2.7	2.5	0.5
1455	Amorpha-4,11-diene	---	---	0.1
1455	α -Humulene	0.6	0.6	0.9
1458	α -Acoradiene	1.8	1.8	---
1461	Alloaromadendrene	2.6	2.5	2.8
1468	β -Acoradiene	4.5	4.6	---
1472	<i>cis</i> -Cadina-1(6),4-diene	0.3	0.3	0.1
1476	<i>trans</i> -Cadina-1(6),4-diene	4.1	4.2	0.5
1478	γ -Muurolole	---	---	1.0
1479	α -Amorphene	---	---	0.9
1480	Germacrene D	---	---	4.0
1481	<i>ar</i> -Curcumene	1.2	1.4	---
1483	(<i>Z,Z</i>)- α -Farnesene	0.4	0.4	0.1
1485	δ -Selinene	---	---	0.2
1487	β -Selinene	---	---	0.4
1490	<i>trans</i> -Muurolo-4(14),5-diene	---	---	0.6
1491	Viridiflorene (Ledene)	1.3	1.3	---

1494	α -Selinene	---	---	0.9
1495	β -Alaskene	3.0	2.9	---
1498	α -Muurolene	0.9	0.9	2.1
1501	β -Cadinene	---	---	0.4
1511	δ -Amorphene	---	---	1.6
1514	β -Curcumene	13.0	12.8	2.2
1522	<i>trans</i> -Calamenene	---	---	0.3
1525	δ -Cadinene	26.0	26.3	53.2
1528	Zonarene	---	---	0.9
1530	(<i>E</i>)- γ -Bisabolene	---	---	0.1
1534	<i>trans</i> -Cadine-1,4-diene	0.1	0.1	0.3
1538	α -Cadinene	0.3	0.3	0.6
1543	α -Calacorene	0.1	0.1	0.7
1546	Elemol	---	---	0.4
1557	Germacrene B	---	---	1.4
1561	(<i>E</i>)-Nerolidol	1.2	1.1	---
1563	β -Calacorene	0.1	0.1	0.2
1571	Maaliol	0.3	0.2	---
1577	Spathulenol	0.1	0.1	tr
1580	<i>trans</i> -Sesquisabinene hydrate	tr	tr	---
1585	Gleenol	---	---	0.1
1587	Globulol	0.8	0.8	0.2
1595	Viridiflorol	0.4	0.4	0.2
1597	<i>allo</i> -Cedrol	0.2	0.2	---
1606	Ledol	0.6	0.6	0.1
1608	Rosifoliol	0.1	0.1	---
1612	Cedrol	2.4	2.5	---
1615	1,10 di- <i>epi</i> -Cubenol	0.3	0.3	0.1
1620	Junenol	---	---	0.1
1621	Selina-6-en-4 β -ol	---	---	0.1
1625	<i>epi</i> -Cedrol	0.5	0.4	---
1628	1- <i>epi</i> -Cubenol	0.5	0.5	1.6
1629	γ -Eudesmol	---	---	0.1
1632	<i>cis</i> -Cadin-4-en-7-ol	---	---	0.2
1633	α -Acorenol	0.2	0.2	---
1638	β -Acorenol	0.1	0.1	---
1644	τ -Cadinol	2.6	2.6	1.3
1647	τ -Muurolol	2.2	2.2	1.8
1648	α -Muurolol (Torreyol)	1.0	1.0	0.8
1659	α -Cadinol	4.7	5.0	2.7
1662	Selin-11-en-4 α -ol	0.1	0.2	0.2
1666	<i>trans</i> -Calamenen-10-ol	tr	tr	0.1
1670	β -Bisabolol	1.2	1.3	tr
1674	Sesquiterpineol	0.1	0.1	---
1685	α -Bisabolol	0.3	0.3	---
1695	Eudesm-7(11)-en-4-ol (Juniper camphor)	---	---	0.1
1723	(<i>Z</i>)-Nuciferol	0.1	0.1	---
1753	β -(<i>Z</i>)-Curcumen-12-ol	0.1	0.1	---
1963	Sandaracopimara-8(14),15-diene	0.1	0.1	---
2132	Nezukol	0.1	tr	---
	Monoterpene hydrocarbons	0.1	0.1	3.4
	Oxygenated monoterpenoids	tr	tr	0.4
	Sesquiterpene hydrocarbons	79.6	79.3	83.9
	Oxygenated sesquiterpenoids	19.9	20.3	10.0
	Diterpenoids	0.2	0.1	0.0
	Total Identified	99.8	99.8	97.6

^a RI = Retention Indices determined with respect to a homologous series of *n*-alkanes on a ZB-5 column.

^b tr = trace (< 0.05%).

The differences in essential oil compositions of *C. odorata* are not surprising. Significant genetic differentiation has been observed between northern Pacific and Atlantic populations of *C. odorata* in Costa Rica [12]. Likewise, there are significant genetic differences between populations of *C. odorata* from the Yucatán of Mexico and those from northwestern Costa Rica and Nicaragua [13]. Not surprisingly, there are also striking differences between the leaf essential oils of *C. odorata* from Nigeria [5], with α -santalene (9.5%), β -

acoradiene (7.1%), β -elemene (6.8%), and caryophyllene oxide (6.0%), as the major components, and the leaf oil from Viçosa, Brazil [6], which had germacrene A (22.6%), β -elemene (19.3%), and bicyclogermacrene (7.6%) as the main essential oil compounds. Similarly, the bark essential oil of *C. odorata* from São Tomé y Príncipe was rich in α -copaene (14.4%), α -cadinol (11.2%), β -eudesmol (9.4%), and δ -cadinene (9.2%) [7]. *Cedrela odorata* bark oil from Costa Rica, on the other hand, was dominated by β -elemene (20.3%)

and germacrene D (15.4%). Although the compositions of *C. odorata* essential oils are remarkably different, they are all dominated by sesquiterpene hydrocarbons regardless of the plant tissue or geographical location.

4. Conclusions

Cedrela odorata wood essential oil has a pleasant aroma and if overharvesting can be prevented, may be an important source of natural ingredients for the cosmetic and flavor industries. The chemical compositions of *C. odorata* wood essential oils are generally dominated by δ -cadinene, but the relative concentrations of minor components vary widely. The organoleptic properties of *C. odorata* wood oil, then, can be profoundly influenced by the geographical source of the oil; a comprehensive analysis of essential oils of *C. odorata* from different geographical locations would be a welcome addition to our knowledge of the phytochemical properties of this tree.

5. Acknowledgments

We are grateful to Leonardo Vargas Garcia, La Cruz, Abangares, Guanacaste, Costa Rica, for providing us with *C. odorata* wood shavings from his workshop. We also thank Daniel Sheffield (Essential Oil Quality Alliance, Charlestown, Indiana, USA) for the sample of *C. odorata* wood essential oil from Colombia. This work was carried out as part of the activities of the Aromatic Plant Research Center (APRC, <https://aromaticplant.org/>). The authors are grateful to dō TERRA International (<https://www.doterra.com/US/en>) for financial support of the APRC.

6. Conflicts of Interest

The authors declare no conflicts of interest. The funding sponsor, dō TERRA International, played no role in the design of the study; in the collection, analysis, or interpretation of the data; conclusions of the study; or in the decision to publish the results.

7. References

1. Morton JF. Atlas of Medicinal Plants of Middle America, Charles C. Thomas, Springfield, Illinois, USA, 1981, I.
2. Cintron BB, *Cedrela odorata* L. In Silvics of North America. Hardwoods. Burns RM, Honkala BH, Eds., Forest Service, United States Department of Agriculture: Washington, DC, USA. 1990; 2:250-257.
3. Ricart Pujals JLR, Padrón Vélez R, Sinopsis Anotada Y. Comentada de la Flora del Bosque Estatal de Maricao. Forest Service, United States Department of Agriculture, Washington, DC, USA, 2010.
4. Lamb AFA. *Cedrela odorata*; University of Oxford, Oxford, UK, 1968.
5. Asekun OT, Ekundayo O. Constituents of the leaf essential oil of *Cedrela odorata* L. from Nigeria. Flavour and Fragrance Journal. 1999; 14:390-392.
6. Maia BHLNS, De Paula JR, Sant Ana J, Silva MF, das GF, Fernandes JB. Essential oils of *Toona* and *Cedrela* species (Meliaceae): Taxonomic and ecological implications. Journal of the Brazilian Chemical Society, 2000; 11:29-639.
7. Martins AP, Salgueiro LR, da Cunha AP, Vila R, Cañigueral S, Tomi F, Casanova J. Chemical composition of the bark oil of *Cedrela odorata* from S. Tomé and Príncipe. Journal of Essential Oil Research. 2003; 15:422-424.
8. Villanueva HE, Tuten JA, Haber WA, Setzer WN. Chemical composition and antimicrobial activity of the

9. Hardt IH, Rieck A, Fricke C, König WA. Enantiomeric composition of sesquiterpene hydrocarbons of the essential oil of *Cedrela odorata* L. Flavour and Fragrance Journal. 1995; 10:16-171.
10. Adams RP. Identification of Essential Oil Components by Gas Chromatography / Mass Spectrometry, 4th ed. Allured Publishing, Carol Stream, Illinois, 2007.
11. Satyal P. Development of GC-MS Database of Essential Oil Components by the Analysis of Natural Essential Oils and Synthetic Compounds and Discovery of Biologically Active Novel Chemotypes in Essential Oils. Ph.D. Dissertation, University of Alabama in Huntsville, 2015.
12. Gillies ACM, Cornelius JP, Newton AC, Navarro C, Hernandez M, Wilson J. Genetic variation in Costa Rican populations of the tropical timber species *Cedrela odorata* L., assessed using RAPDs. Molecular Ecology. 1997; 6:1133-1145.
13. Cavers S, Navarro C, Lowe AJ. Chloroplast DNA phylogeography reveals colonization history of a neotropical tree, *Cedrela odorata* L., in Mesoamerica. Molecular Ecology. 2003; 12:1451-1460.